Pharmacy Joe's Critical Care Pharmacy Academy Masterclass (pharmacyjoe.com)

ICU Rounds: Before, During, and After

- 1 Objectives
  - 1.1 List the essential health care practitioners that should be available on ICU rounds
  - 1.2 Outline a process to use before, during, and after rounds
  - 1.3 Select parts of the rounding processes described to incorporate into daily practice
- 2 Before Rounds a.k.a "Pre-Rounding"
  - 2.1 Patient Observation
    - 2.1.1 Do pre-rounding in front of the patient when possible
    - 2.1.2 How sick do they look?
    - 2.1.3 Intubated?
    - 2.1.4 SCDs on?
    - 2.1.5 Tube feedings active?
    - 2.1.6 Better collaboration with nurse or physician
  - 2.2 Rounding Tools
    - 2.2.1 General philosophy
    - 2.2.2 What I write on my rounding tool
    - 2.2.3 Room, Name, Wt, Age, Gender
    - 2.2.4 CICU
    - 2.2.5 Reason for admission
    - 2.2.6 VAP
    - 2.2.7 SUP
      - 2.2.7.1 If expected to be ventilated >24 hours
      - 2.2.7.2 Coagulopathy (INR >2 or aPTT >45s)
      - 2.2.7.3 Full-dose anticoagulation
      - 2.2.7.4 Vasopressor dose >10 mcg/min
      - 2.2.7.5 NSAID or glucocorticoid use
      - 2.2.7.6 Mechanical ventilation > 24 hours
      - 2.2.7.7 Multi-system organ failure
      - 2.2.7.8 History of peptic ulcer disease
      - 2.2.7.9 Acute renal failure
      - 2.2.7.10 H2RA vs PPI
      - 2.2.7.11 Enteral nutrition as prophylaxis?
      - 2.2.7.12 Evaluate for discontinuation daily and ESPECIALLY on transfer from ICU
      - 2.2.7.13 ??
    - 2.2.8 VTE
      - 2.2.8.1 All ICU patients are at high risk of VTE
      - 2.2.8.2 SCDs
      - 2.2.8.3 Low dose heparin or enoxaparin
      - 2.2.8.4 Platelets must be at least 50,000
      - 2.2.8.5 Heparin bid vs tid
      - 2.2.8.6 Epidurals
      - 2.2.8.7 Check with your anesthesia department for local guidelines
      - 2.2.8.8.1 ASRA recommends a minimum of 2 hours between placement and VTE prophylaxis. FDA recommends 4 hours
      - 2.2.8.9 Antibiotic days
      - 2.2.8.10 Therapeutic drug monitoring (heparin, vancomycin, aminoglycosides, phenytoin, etc...)
      - 2.2.8.11 Recommendations
      - 2.2.8.12 Pharmacy Problem List
      - 2.2.8.13 RICHMAN
        - 2.2.10.1.1 Respiratory
        - 2.2.2.10.1.2 Infectious
        - 2.2.2.10.1.3 Cardiac
        - 2.2.2.10.1.4 Hematologic
        - 2.2.2.10.1.5 Metabolic
        - 2.2.2.10.1.6 Alimentary
        - 2.2.2.10.1.7 Neurologic
      - 2.2.2.10.1.8 Alternatives (head-to-toe)
      - 2.2.2.10.2.1 An alternative tool: FASTHUG MADDINS
    - 2.2.9 Feeding
    - 2.2.9.1.1 Analgesia
    - 2.2.9.1.2 Sedation
    - 2.2.9.1.3 Thromboprophylaxis
    - 2.2.9.1.4 Hypoactive or Hyperactive delirium (sometimes Head of bed elevation)
    - 2.2.9.1.5 stress Ulcer prophylaxis
    - 2.2.9.1.6 Glucose control
    - 2.2.9.1.7 Medication reconciliation
    - 2.2.9.1.8 Antibiotics
    - 2.2.9.1.9 Indications for medications
    - 2.2.9.1.10 Drug dosing
    - 2.2.9.1.11 Electrolytes, hematology, and other laboratory tests
    - 2.2.9.1.12 No drug interactions, allergies, duplication, or side effects
    - 2.2.9.1.13 Stop dates
    - 2.2.9.1.14 5 Step pre-rounding process
  - 2.3 1. Who is the patient and why are they here?
    - 2.3.1 Focus on the History of present illness & Assessment/Plan of the ICU physician
    - 2.3.1.1 Other documents to review
    - 2.3.2 History & Physicial
    - 2.3.2.1 Consult notes
    - 2.3.2.2 Progress notes
    - 2.3.2.3 Nursing notes
    - 2.3.2.4 2. Review home medications
    - 2.3.3 Is Medication Reconciliation complete?
    - 2.3.3.1 Will holding home meds cause a problem?
    - 2.3.3.2 Beta blockers
    - 2.3.3.2.1 Clonidine
    - 2.3.3.2.2 Opioids
    - 2.3.3.2.3 Benzodiazepines
    - 2.3.3.2.4 Antidepressants
    - 2.3.3.2.5 Anti-rejection
    - 2.3.3.2.6 Anti-retroviral
    - 2.3.3.2.7 Could home meds have caused admission?
    - 2.3.3.3 3. Review Additional Data
    - 2.3.4 Labs
    - 2.3.4.1 Cultures
    - 2.3.4.2 Imaging
    - 2.3.4.3 EKG
    - 2.3.4.4 4. Review Bedside Flowsheet
    - 2.3.5 Respiratory Requirements
      - 2.3.5.1 FiO2
        - 2.3.5.1.1 40% is minimum
        - 2.3.5.1.1.1 Mode of oxygenation
        - 2.3.5.1.2 Room air
        - 2.3.5.1.2.1 Nasal Canula
        - 2.3.5.1.2.2 BiPAP
        - 2.3.5.1.2.3 Endotracheal
        - 2.3.5.1.2.4 PEEP
        - 2.3.5.1.2.5 5 is minimum
        - 2.3.5.1.2.5.1 Sedation Scale
        - 2.3.5.2 Glasgow Score
        - 2.3.5.3 Vital signs
        - 2.3.5.4 Current infusions
        - 2.3.5.5 Dosing & number of sedatives vs sedation score
        - 2.3.5.5.1 Dosing & number of vasopressors vs vital signs
        - 2.3.5.5.2 Input & Output
        - 2.3.5.6 At least 0.5 mL/kg/min urine output
        - 2.3.5.6.1 Oliguria
        - 2.3.5.6.2 Anuria
        - 2.3.5.6.3 5. Review current medications
    - 2.3.6 Do this last to avoid the guessing game trap
    - 2.3.6.1 It is easier to recognize medication related problems if you have evaluated everything else first
    - 2.3.7 SCr normal but no urine output
    - 2.3.7.1 Agitated patient on ventilation at home
    - 2.3.7.2 Septic shock from abdominal source but only on levofloxacin/metronidazole
    - 2.3.7.3 Difficult to sedate patient on buprenorphine at home
    - 2.3.7.4 Propofol only (no analgesia) in a patient with significant surgery
    - 2.3.7.5 Who to see first?
  - 2.4 Quickly walk the unit before starting to find the sickest patient(s)
    - 2.4.1 On CRRT
    - 2.4.1.1 Numerous IV pumps running
    - 2.4.1.2 Code cart pre-emptively placed outside room by nursing staff
    - 2.4.1.3 Collaborate with the charge nurse "Who is the sickest patient in the unit?"
    - 2.4.1.4 How long to spend per patient?
    - 2.4.2 20-30 minutes for a new patient
    - 2.4.2.1 10-15 minutes for an existing patient
    - 2.4.2.2 Utilize residents and students as "pharmacist extenders"
    - 2.4.2.3 HAVE AN ABBREVIATED PROCESS
    - 2.5.4
- 3 During Rounds
  - 3.1 Strongly recommended Best Practices from Critical Care Medicine 2013 article:
    - 3.1.1 A Systematic Review of Evidence-Informed Practices for Patient Care Rounds in the ICU
    - 3.1.1 Implement multidisciplinary rounds (including at least a medical doctor, registered nurse, and pharmacist)
    - 3.1.1.1 Standardize location, time, and team composition
    - 3.1.2 Define explicit roles for each HCP participating on rounds
    - 3.1.3 Develop and implement structured tool (best practices checklist)
    - 3.1.4 Reduce nonessential time wasting activities
    - 3.1.5 Minimize unnecessary interruptions
    - 3.1.6 Focus discussions on development of daily goals and document all discussed goals in health record
    - 3.1.7 Structure of my ICU rounds
  - 3.2 Team
    - 3.2.1 Physician
    - 3.2.1.1 Bedside nurse
    - 3.2.1.2 Pharmacist
    - 3.2.1.3 Charge nurse
    - 3.2.1.4 Respiratory therapist
    - 3.2.1.5 Dietician
    - 3.2.1.6 Spiritual care
    - 3.2.1.7 Location - outside of room 1
    - 3.2.2 Time - 10am
    - 3.2.3 Rules for each discipline
    - 3.2.4 Physician - briefly presents the patient
    - 3.2.4.1 Bedside nurse - describes overnight events
    - 3.2.4.2 Physician or nurse - review systems using RICHMAN
    - 3.2.4.3 Respiratory
    - 3.2.4.3.1 Infectious
    - 3.2.4.3.2 Cardiac
    - 3.2.4.3.3 Hematologic
    - 3.2.4.3.4 Neurologic
    - 3.2.4.3.5 Alimentary
    - 3.2.4.3.6 Metabolic
    - 3.2.4.3.7 Each discipline is expected to offer any recommendations or ask any appropriate questions as the systems are reviewed and before moving to the next patient
    - 3.2.4.4 Role for the pharmacist on rounds
  - 3.3 Tailor your role to the unique circumstances of your practice
    - 3.3.1 Nutrition services present?
    - 3.3.1.1 Computer system ensures medication renewals are addressed?
    - 3.3.1.2 Head of bed elevation consistently evaluated by other practitioners?
    - 3.3.1.3 Monitor all medication use
    - 3.3.2 Safety
    - 3.3.2.1 Toxicity
    - 3.3.2.1.1 Allergy
    - 3.3.2.1.2 Drug interactions
    - 3.3.2.1.3 Adverse drug events
    - 3.3.2.1.4 Renal / hepatic dose adjustment
    - 3.3.2.1.5 Efficacy
    - 3.3.2.2 Therapeutic drug monitoring when possible (vancomycin, aminoglycosides, phenytoin, etc...)
    - 3.3.2.2.1 Cost-effectiveness / stewardship
    - 3.3.2.3 Provide drug information
    - 3.3.3 Compatibility
    - 3.3.3.1 Monitoring guidelines
    - 3.3.3.2 Answer specific questions
    - 3.3.3.3 Provide education for other disciplines
    - 3.3.4 When and how to present your recommendations
  - 3.4 The most effective time to give recommendations to a provider is when they are trying to solve the patient's problems
    - 3.4.1 Be elbow-to-elbow with the providers as they are "seeing the patient" and determining orders
    - 3.4.1.1 Getting the best recommendation accepted at the beginning of care is so much easier than trying to change therapy later
    - 3.4.1.2 Present "basic" recommendations like prophylaxis issues outside of rounds
    - 3.4.2 Place an 'unofficial' note in a spot that you know is within a physician's workflow (I use a sticky note on the nurse's flowsheet)
    - 3.4.2.1 Why?
    - 3.4.2.2 Saves rounding time
    - 3.4.2.2.1 Avoids "embarrassing" the MD about something simple that was overlooked
    - 3.4.2.2.2 Allows you to double check during rounds that the recommendation was accepted
    - 3.4.2.2.3 During rounds, make recommendations when the relevant system is being discussed
    - 3.4.3 Stress ulcer recommendations during "Alimentary"
    - 3.4.3.1 Antibiotic changes during "Infectious"
    - 3.4.3.2 Diffuse potentially adversarial situations
    - 3.4.4 Has the physician "seen" the patient yet?
    - 3.4.4.1 "I know you haven't seen the patient yet..."
    - 3.4.4.1.1 What are the goals of care (curative vs palliative)?
    - 3.4.4.2 Pharmacist as a "blind golf caddy"
    - 3.4.4.2.1 For controlling agitation I ask: Can we intubate this patient if we give too many sedatives? This helps me decide which sedatives I should use and which I should avoid.
    - 3.4.4.2.1.1 For weaning off long-term fentanyl infusions I ask: Is the fentanyl infusion the only thing keeping the patient in the ICU? This helps me choose how aggressively to wean the fentanyl.
    - 3.4.4.2.1.2 Do you think the patient might have necrotizing fasciitis? Helps decide whether to add a dose of clindamycin to decrease toxin production before the patient goes to the OR for source control.
    - 3.4.4.2.1.3 How to approach recommendations when care is futile
    - 3.4.4.2.1.4
- 4 After Rounds
  - 4.1 Prioritize which problems to solve first
  - 4.1.1 Have a system to ensure that tasks are recorded & completed
  - 4.2 Reminders
  - 4.2.1 Reviewing pre-rounding tools
  - 4.2.2 Follow-up as needed
  - 4.3 Did the physician change orders as discussed?
  - 4.3.1 Does the bedside nurse need additional education on how to use or monitor a medication?
  - 4.3.2 Monitor for safety / efficacy
  - 4.3.3 TDM
  - 4.3.4 Evaluate patients further if pre-rounding time was limited
  - 4.4